## PH481 Homework 8

Due: Friday, 6th of March 2020

**8.4** Describe completely the state of polarization of each of the following waves:

(a) 
$$\vec{\mathbf{E}} = \hat{\mathbf{i}} E_0 \cos(kz - \omega t) - \hat{\mathbf{j}} E_0 \cos(kz - \omega t)$$

(b) 
$$\vec{\mathbf{E}} = \hat{\mathbf{i}}E_0 \sin 2\pi (z/\lambda - \nu t) - \hat{\mathbf{j}}E_0 \sin 2\pi (z/\lambda - \nu t)$$

(c) 
$$\vec{\mathbf{E}} = \hat{\mathbf{i}}E_0 \sin(\omega t - kz) + \hat{\mathbf{j}}E_0 \sin(\omega t - kz - \pi/4)$$

(d) 
$$\vec{\mathbf{E}} = \hat{\mathbf{i}}E_0 \cos(\omega t - kz) + \hat{\mathbf{j}}E_0 \cos(\omega t - kz + \pi/2)$$
.

**8.8\*** Write an expression for a  $\mathcal{P}$ -state lightwave of angular frequency  $\omega$  and amplitude  $E_0$  propagating along a line in the xy-plane at 45 ° to the x-axis and having its plane-of-vibration corresponding to the xy-plane. At t=0, x=0, and y=0 the field is zero.

**8.26\*** Imagine a pair of crossed polarizers with transmission axes vertical and horizontal. The beam emerging from the first polarizer has flux density  $I_1$ , and of course no light passes through the analyzer (i.e.,  $I_2 = 0$ ). Now insert a perfect linear polarizer (*HN*-50) with its transmission axis at 45° to the vertical between the two elements—compute  $I_2$ . Think about the motion of the electrons that are radiating in each polarizer.

**8.27\*** Imagine that you have two identical perfect linear polarizers and a source of natural light. Place them one behind the other and position their transmission axes at  $0^{\circ}$  and  $50^{\circ}$ , respectively. Now insert between them a third linear polarizer with its transmission axes at  $25^{\circ}$ . If  $_{1000}$  W/m<sup>2</sup> of light is incident, how much light will emerge with and without the middle polarizer in place?

## HW 8 extra problems

- A. Consider a plane wave with wavelength  $\lambda$  incident normally on a screen with a circular aperture of radius a. The point of observation is directly opposite the center of the aperture at a distance of  $r_0 = 2a^2/\lambda$  from the screen  $(r_0 \gg \lambda)$ .
  - a) How many Fresnel zones are contained in the aperture as seen from the observation point?
  - b) Draw the vibration curve and the phasor corresponding to this case.
  - c) What is the intensity at the observation point in terms of the intensity with the screen absent?
- B. Draw the Cornu spiral. Consider a long slit that contains one Fresnel zone. Discuss and show how you would use the Cornu spiral to find the intensity at a point directly opposite the slit. How does this intensity compare with that from a slit that contains two Fresnel zones?

Other PRACTICE problems (no need to turn in; will not be graded) Hecht: 10.72, 10.74, 10.76, 10.81, 10.84, 10.92